

University of Groningen

Identifying markers for premature atherosclerosis in rheumatoid arthritis

de Groot, Lodewijk

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

de Groot, L. (2016). *Identifying markers for premature atherosclerosis in rheumatoid arthritis*. [Thesis fully internal (DIV), University of Groningen]. Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHAPTER

Introduction

1

1

General Introduction

Cardiovascular morbidity and mortality are increased in rheumatoid arthritis (RA) (1-3). Despite improved treatment of RA an increased incidence of cardiovascular diseases still remains, as is demonstrated by recent large cohort studies (4-6). Various causes of this increased prevalence have been proposed. Of these, traditional risk factors like smoking, hypertension, hyperlipidemia and increased body mass index (BMI) are involved (7-9). Also non traditional risk factors like oxidized LDL (ox-LDL), advanced glycation end products (AGEs) and auto-antibodies play a role (10).

Epidemiology

An increased risk for cardiovascular morbidity and mortality in RA has been shown. The nurses health study showed a twofold increase in myocardial infarction in women with RA, increasing to more than threefold when disease duration extended longer than ten years (11). In a large meta-analysis an increased mortality of 50% was found due to cardiovascular disease (CVD) in RA (1). Risk for myocardial infarction proved to be greater than that in type 2 diabetes mellitus (T2DM) in one study (12).

Markers for identifying premature atherosclerosis

Various markers are used for identifying premature atherosclerosis. Different stages of atherosclerosis are characterized by specific markers. These are extensively discussed in chapter two. The first stage is characterized by invasion of inflammatory cells into the vascular wall. Markers of this stage are, for example, serum levels of vascular cellular adhesion molecule-1 (sVCAM-1), von Willebrand factor (vWF) and thrombomodulin (TM). The second stage is characterized by endothelial dysfunction through further inflammation of the vascular wall. This leads to "stiffness" of the arterial wall, which is characterized by increased velocity of the pulse wave from the brachial artery to the radial artery. This increased velocity can be visualized by pulse wave analysis (PWA)(13). The increase in velocity is recalculated by a windkessel model to small artery elasticity (SAE). Finally the process leads to thickening of the vascular wall, as measured by intima media thickness (IMT). IMT is measured on the left and right side of the carotid bulb (CB), the internal carotid artery (ICA) and the common carotid artery (CCA). Increment of IMT in RA has been shown in several studies as compared to healthy controls (14, 15).

Aim and outline of the thesis

The main aim of the present thesis is (1) to identify markers for detection of early atherosclerosis in RA, and (2) to evaluate the effect of disease activity and treatment on early markers of atherosclerosis.

Introduction of the chapters

Chapter 2 discusses epidemiology and pathophysiology of cardiovascular disease in RA. Also methods for investigating the several stages of preclinical cardiovascular disease are discussed, such as markers for endothelial activation, endothelial dysfunction, IMT and advanced glycation endproducts (AGE's). Therapeutic strategies for minimizing cardiovascular risk are discussed in this chapter as well. This invited review (allowing limited number of references) was published in 2010 so covering the literature until that time.

Chapter 3 discusses a cross sectional study in 49 RA patients with established disease. We focused in this study on SAE. We hypothesized that the decrease in SAE in RA would be related to disease activity and markers of disease damage like the Sharp v.d. Heijde score, as well as to markers of endothelial activation and IMT.

In chapter 4 we cross-sectionally studied 49 RA patients with longstanding disease in comparison with sex and age matched healthy controls for the presence of AGE's. AGE's are the result of non-enzymatic oxidation and glycation of proteins and lipids under a hyperglycemic state or oxidative stress. Patients with long standing RA are supposed to have had longer periods of oxidative stress due to their disease. Here we related AGE's to markers of endothelial activation, such as sVCAM-1 and vWF, endothelial dysfunction as measured by small artery elasticity (SAE) and premature atherosclerosis as measured by IMT.

We hypothesized that in longstanding RA, AGE's are increased due to prolonged exposure to oxidative stress and as such, correlate with markers of disease damage. Furthermore we hypothesized that AGE accumulation is related to endothelial activation, endothelial dysfunction, and early atherosclerosis.

In chapter 5 we focus on angiogenic factors (angiopoietin-2 (Angpt-2) and vascular endothelial growth factor (VEGF)) in a cohort of 176 RA patients with established disease. This cohort was followed for 12.5 years and prevalence of CVD was retrieved from medical records.

As angiogenesis plays an important role in inflammation and, as such, possibly also in the development of atherosclerosis we analyzed angiogenic factors in relation to atherosclerosis in RA. Whether angiogenesis is a leading factor in inflammation or secondary to hypoxia in inflamed tissue is, however, still under debate (16) We here hypothesized that angiogenic factors would correlate with the development of CVD. Angpt-2, as well as VEGF and sVCAM-1, were measured at entry and after 2 years and were compared with levels in healthy controls (HC).

Chapter 6 concerns the role of endothelial progenitor cells (EPC's). EPC's are thought to play an important role in endothelial repair (17, 18) Depletion of EPC's might be related to defective endothelial repair, which could play a role in atherosclerosis. We studied an early RA cohort of 27 RA patients. EPC's were defined as CD133/CD34 positive cells by

1

FACS analysis or by colony forming units by the Hill assay (19, 20). Numbers and function of EPC's were compared with that in age and sex matched healthy controls. Correlation with markers of endothelial activation and dysfunction as well as disease activity were studied.

In chapter 7 a longitudinal study is described in which 58 RA patients with a maximum disease duration of one year were treated with standard treatment consisting mainly of methotrexate, sulfasalazine and TNF α -blocking agents. We hypothesized that markers of endothelial dysfunction would be present before start of treatment and that disease activity would correlate with markers of early atherosclerosis. Also, we hypothesized that by reducing disease activity in the first year of treatment markers of premature atherosclerosis would decrease. Primary outcome in this study was endothelial function as measured by SAE. SAE was measured before initiation of therapy and after one year. Results found at entry were compared with that of age and sex matched healthy controls.

Secondary outcome measures were sVCAM-1 as an endothelial activation marker, intima media thickness and advanced glycation end products (AGEs). As mentioned premature markers for atherosclerosis were measured before start of treatment and after one year.

Finally, chapter 8 discusses a pilot longitudinal study in 15 RA patients with longstanding disease, who were treated with rituximab. We hypothesized that before rituximab treatment markers of premature cardiovascular disease would be present, and correlate with disease activity and that markers of premature cardiovascular disease would decrease after treatment. Primary outcome in this study was endothelial activation, as measured by sVCAM-1. Secondary outcome measures were endothelial dysfunction and IMT. Indication for treatment with rituximab in these patients was established by their own rheumatologist. After giving informed consent patients were included in the study. Before rituximab treatment markers of early cardiovascular disease were measured and compared with that of age and sex matched healthy controls. Four months after rituximab treatment measurements were repeated in the RA group. S-VCAM-1, SAE and IMT were related to disease activity.

All results are discussed in chapter 9 (Summary conclusions and future recommendations)

References

1. Avina-Zubieta JA, Choi HK, Sadatsafavi M, Etminan M, Esdaile JM, Lacaille D. Risk of cardiovascular mortality in patients with rheumatoid arthritis: a meta-analysis of observational studies. *Arthritis Rheum.* 2008 12;15;59(0004-3591; 0004-3591; 12):1690-7.
2. del Rincon ID, Williams K, Stern MP, Freeman GL, Escalante A. High incidence of cardiovascular events in a rheumatoid arthritis cohort not explained by traditional cardiac risk factors. *Arthritis Rheum.* 2001 12;44(0004-3591; 0004-3591; 12):2737-45.
3. Solomon DH, Reed G, Kremer JM, Curtis JR, Farkouh ME, Harrold LR, et al. Disease activity in rheumatoid arthritis and the risk of cardiovascular events. *Arthritis Rheumatol.* 2015 Mar 16.
4. Sparks JA, Chang SC, Liao KP, Lu B, Fine AR, Solomon DH, et al. Rheumatoid arthritis and mortality among women during 36 years of prospective follow-up: Results from the Nurses' Health Study. *Arthritis Care Res (Hoboken).* 2015 Oct 16.
5. Van Doornum S, Bohnensky M, Tacey MA, Brand CA, Sundararajan V, Wicks IP. Increased 30-day and 1-year mortality rates and lower coronary revascularisation rates following acute myocardial infarction in patients with autoimmune rheumatic disease. *Arthritis Res Ther.* 2015 Feb 27;17:38,015-0552-2.
6. Mackey RH, Kuller LH, Deane KD, Walitt BT, Chang YF, Holers VM, et al. Rheumatoid Arthritis, Anti-Cyclic Citrullinated Peptide Positivity, and Cardiovascular Disease Risk in the Women's Health Initiative. *Arthritis Rheumatol.* 2015 Sep;67(9):2311-22.
7. Kremers HM, Crowson CS, Therneau TM, Roger VL, Gabriel SE. High ten-year risk of cardiovascular disease in newly diagnosed rheumatoid arthritis patients: a population-based cohort study. *Arthritis Rheum.* 2008 08;58(0004-3591; 8):2268-74.
8. Lewington S, Whitlock G, Clarke R, Sherliker P, Emberson J, Halsey J, et al. Blood cholesterol and vascular mortality by age, sex, and blood pressure: a meta-analysis of individual data from 61 prospective studies with 55,000 vascular deaths. *Lancet.* 2007 12/01;370(1474-547; 1474-547; 9602):1829-39.
9. Nurmohamed MT. Cardiovascular risk in rheumatoid arthritis. *Autoimmun Rev.* 2009 07;8(1873-0183; 1568-9972; 8):663-7.
10. Matsumoto T, Tsurumoto T, Baba H, Osaki M, Enomoto H, Yonekura A, et al. Measurement of advanced glycation endproducts in skin of patients with rheumatoid arthritis, osteoarthritis, and dialysis-related spondyloarthropathy using non-invasive methods. *Rheumatol Int.* 2007 12;28(1437-160; 0172-8172; 2):157-60.
11. Solomon DH, Karlson EW, Rimm EB, Cannuscio CC, Mandl LA, Manson JE, et al. Cardiovascular morbidity and mortality in women diagnosed with rheumatoid arthritis. *Circulation.* 2003 Mar 11;107(9):1303-7.
12. van Halm VP, Peters MJ, Voskuyl AE, Boers M, Lems WF, Visser M, et al. Rheumatoid arthritis versus diabetes as a risk factor for cardiovascular disease: a cross-sectional study, the CARRE Investigation. *Ann Rheum Dis.* 2009 09;68(1468-2060; 1468-2060; 9):1395-400.
13. Van Doornum S, McColl G, Jenkins A, Green DJ, Wicks IP. Screening for atherosclerosis in patients with rheumatoid arthritis: comparison of two in vivo tests of vascular function. *Arthritis Rheum.* 2003 01;48(0004-3591; 1):72-80.
14. Gonzalez-Gay MA, Gonzalez-Juanatey C, Vazquez-Rodriguez TR, Martin J, Llorca J. Endothelial Dysfunction, Carotid Intima-Media Thickness, and Accelerated Atherosclerosis in Rheumatoid Arthritis. *Semin Arthritis Rheum.* 2008 04/04(0049-0172).
15. Daza L, Aguirre M, Jimenez M, Herrera R, Bollain JJ. Common carotid intima-media thickness and von Willebrand factor serum levels in rheumatoid arthritis female patients without cardiovascular risk factors. *Clin Rheumatol.* 2007 04;26(0770-3198; 4):533-7.
16. Costa C, Incio J, Soares R. Angiogenesis and chronic inflammation: cause or consequence? *Angiogenesis.* 2007;10(3):149-66.
17. Hristov M, Zerneck A, Liehn EA, Weber C. Regulation of endothelial progenitor cell homing after arterial injury. *Thromb Haemost.* 2007 Aug;98(2):274-7.
18. Rosenzweig A. Circulating endothelial progenitors--cells as biomarkers. *N Engl J Med.* 2005 Sep 8;353(10):1055-7.
19. Asahara T, Murohara T, Sullivan A, Silver M, van der Zee R, Li T, et al. Isolation of putative progenitor endothelial cells for angiogenesis. *Science.* 1997 Feb 14;275(5302):964-7.
20. Hill JM, Zalos G, Halcox JP, Schenke WH, Waclawiw MA, Quyyumi AA, et al. Circulating endothelial progenitor cells, vascular function, and cardiovascular risk. *N Engl J Med.* 2003 Feb 13;348(7):593-600.

